

15/5/20 OK! (but big mistakes...)

IE CHEMISTRY n° 2

Pierre  
Petrella

10,75  
12

I- Partial miscibility at liquid state

A-Liquid/vapor equilibria:

0,5  
3- to obtain pure B by fractionated distillation we would have to have an initial liquid composition of B bigger than 0,4 such that once all the vapor is taken away through heating (of heteroazeotropic proportions) we will only have B left in the liquid state.

25°C 1000 moles of A and B with molar fraction of B = 0,10

0,5  
4). under 1 bar the mixture will start boiling at 97°C (according to the graph)

the composition of the vapor formed initially will be of  $y_B = 0,26$  in molar fractions.  
hence

0,5  
5) the composition of the distillate will be of  $D_1 = 0,4$

0,25  
6) in the residue we will only have A left.  $R_1 = 0$

7) we have initially 900 moles of A and 100 moles of B


in the  $D_1$  we would then have  $\frac{100}{100+x} = 0,4 \Rightarrow 100 = 40 + 94x$   
 $x = 150$

$n_A = 150$  moles and  $n_B = 100$  moles

1,75

in  $R_1$  we would have  $n_A = 750 \text{ mol}$   $n_B = 0$ .

A



B - liquid liquid equilibria  $p = 1 \text{ bar}$

A

- 9) limit of solubility of A in B at  $10^\circ\text{C}$  is  $x_B = 0,96$   
 10) limit of solubility of B in A at  $10^\circ\text{C}$  is  $x_B = 0,10$   
 (found using the table)

11)  $D_1$  is of composition 0,4 we will use the lever rule to find the number of moles in each phase  $n_A$  and  $n_B$ .

we know that  $n_A \cdot \overline{L_1 D_1} = n_B \cdot \overline{D_1 L_2}$

moreover:  $n_A = \frac{\overline{D_1 L_2}}{\overline{L_1 L_2}} n_{\text{total}}$

$$\left. \begin{aligned} n_A &= n_{\text{total}} - n_B \\ n_{\text{total}} \overline{D_1 L_2} - n_B \overline{D_1 L_2} \\ n_A &= \frac{\overline{D_1 L_2}}{\overline{L_1 D_1} + \overline{D_1 L_2}} \end{aligned} \right\}$$

hence  $n_A = \frac{0,96 - 0,4}{0,96 - 0,1} \times 250 = 162,8 \text{ moles}$

and  $n_B = \frac{\overline{L_1 D_1}}{\overline{L_1 L_2}} \times n_{\text{total}} = \frac{0,4 - 0,1}{0,96 - 0,1} = 87,2 \text{ moles}$

(2)

now  $n_A(p)$  has 152,8 moles  
 $n_B(p)$  has 87,2 moles

we get for  $n(p)$   $\begin{cases} n_B = n(p) \times x_B = 16,3 \text{ mol} \\ n_A = n(p) \times (1-x_B) = 146,5 \text{ mol.} \end{cases}$

and  $n(p)$   $\begin{cases} n_B = n(p) \times x_B = 83,7 \text{ mol} \\ n_A = n(p) \times (1-x_B) = 3,5 \text{ mol.} \end{cases}$

C - Second distillation:

12) we take  $B(p)$  solution of  $x_B = 0,96$  and with  $n = 87,2$  moles

same here we will have a distillate of composition of 0,4 in the distillate which will have all of the A moles.

$n_A = 3,5$  moles and  $\frac{n_B}{n_A + n_B} = 0,4 \Rightarrow n_B = 0,4 n_A + 0,4 n_B$   
 $n_A = 4,3$   
 with  $x_B = 0,4 \Rightarrow n_B = \frac{1,4}{0,6} = 2,3 \text{ mol}$

in the remainder,  $\begin{cases} n_A = 0 \\ n_B = 83,7 - 2,3 = 81,4 \end{cases}$   
 with  $x_B = 1$

13)  $100 \times \frac{n_A \text{ recovered pure}}{n_A \text{ initial}} = 100 \times \frac{750}{900} = 83,3$  they are  
 $100 \times \frac{n_B \text{ recovered pure}}{n_B \text{ initial}} = 100 \times \frac{81,4}{100} = 81,4$  quite similar.

4,5

4,75  
8

Exercice n° 2:

1) since  $E^\circ(O_2/H_2O) = 1,23V$  and  $E^\circ(C_6H_{12}O_7/C_6H_{12}O_6) = 0,076V$


0,5

the electrons will travel from the highest to the lowest potential making the highest potential the anode and the lowest potential the cathode.

oxidation will take place at M'

reduction will take place at M

0,5




2)

3) anode:  $E(O_2/H_2O) = E^\circ(O_2/H_2O) + \frac{RT}{nF} \times \ln \left( \frac{[H_3O^+]^4 [O_2]}{1} \right)$


0,75  
cathode.

$E(C_6H_{12}O_7/C_6H_{12}O_6) = E^\circ(C_6H_{12}O_7/C_6H_{12}O_6) + \frac{RT}{nF} \times \ln \left( \frac{[H_3O^+] [C_6H_{12}O_7]}{[C_6H_{12}O_6]} \right)$

4) overall redox reaction:



0,5



5) Find  $K_{298K}$  of this reaction:

since we have a redox reaction we can use that

$$\Delta_r G^\circ = -\Delta E n F = -RT \ln K^\circ$$

$$\Rightarrow \Delta E n F = RT \ln K^\circ$$

$$K^\circ = \exp \left( \frac{\Delta E n F}{RT} \right) = \exp \left( \frac{1,16 \times 4 \times 96500}{8,314 \times 298} \right)$$

A

$$= E(O_2/H_2O) - E(C_6H_{12}O_7/C_6H_{12}O_6) \quad K^\circ = 3,076 \times 10^{42}$$

3,25

1. Pierre  
 2. Kitzells  
 71. 0,5

I find that  $k_i$  is too high to be true, but if it is correct, it is over  $10^4$  which makes the reaction complete in the direct direction. ~~yes~~

electromotive force is around 0,8V.

b) for 1 mole of glucose.

$$E(\text{O}_2/\text{H}_2\text{O}) - E(\text{C}_6\text{H}_{12}\text{O}_6/\text{C}_6\text{H}_{12}\text{O}_5) = 0,8 \text{ V}$$

1 mol will give out a total of ...

$$W_{\text{elec}} = -0,8 \times 4 \times 96500 \times 1 = -308800 \text{ J}$$

of  
 for 1 liter of blood we have 1g of glucose hence

$$n = \frac{1}{180} = 5,56 \times 10^{-3} \text{ mol.}$$

$$W_{\text{elec}}' = -0,8 \times 4 \times 96500 \times 5,56 \times 10^{-3} = 1716,928$$

7) the life span would be  $\frac{W(\text{opt with 1L of blood})}{W = (xU)}$  in s.

$$\frac{W_{\text{elec}}'}{10 \times 10^{-6}} = 171 \times 10^6 \text{ s} \rightarrow 5,4 \text{ years.}$$

05

2) advantages: we won't have to carry an external  
battery.  
disadvantages, it doesn't last forever.

literal expression:  $V = N - r - r' + n - \gamma$




Figure 1 : Isobaric L/V equilibrium phase diagram of A-B binary system under  $P = 1$  bar.

Exercise I - A - 2. Preliminary questions. Check the statements that you think are correct

At the liquid state, A and B mixtures can be considered as :

- Ideal solutions
- Non-ideal solutions

At the liquid state, compared to A-A and B-B molecular interactions, A-B molecular interactions are :

- Of the same order of magnitude
- Stronger
- Weaker

since more volatile.




Figure 2 : Schematic representation of a glucose / dioxygen biological battery cell.